

4. Ordered response models

4.1 General model approaches

Ordered (ordinal) dependent variables in a microeconomic analysis:

These qualitative variables have also more than two possible mutually exclusive categories which are (in contrast to multinomial variables), however, naturally (rank) ordered (in the case of two categories, the variables are binary)

Examples for microeconomic analyses with ordered response models:

- Analysis of the individual satisfaction of a person with life (e.g. on an eleven-point scale of integers from zero for “completely dissatisfied” to ten for “completely satisfied”)
- Analysis of the personal strength of agreement to a political program (e.g. with the categories “strong disagreement”, “weak disagreement”, “weak agreement”, “strong agreement”)
- Analysis of the years of education of a person (e.g. with the categories “less than nine years”, “between nine and 12 years”, “at least 13 years”)
- Analysis of the credit rating of firms (e.g. on a scale from D to AAA)
- Analysis of the stated importance of equity issues in international climate negotiations (e.g. on a five-point scale from “no importance” to “very high importance”)

Preliminary remarks to ordered dependent variables:

- The values for the categories of the ordered dependent variables are completely arbitrary if they preserve the rank order so that e.g. the sequences 1, 2, 3, 4 or 10, 20, 30, 40 or -20, -10, 0, 10 are possible and all reveal the same information for an ordinal variable with four categories (e.g. “strong agreement”, “weak agreement”, “weak disagreement”, “strong disagreement”). As a consequence, expectations, variances, or covariances for values of ordinal variables have no meaning.
- For notational simplicity, the dependent ordered variables y_i of an observation i take the values 1, 2, ..., J for the categories where “1” < “2” < ... < “ J ”
- In general, ordered dependent variables could therefore be analyzed with multinomial response models as discussed before. However, this ignores the ordering information so that this would lead to inefficient ML estimators of the parameters compared with the use of ordered response models.
- This coding 1, 2, ..., J should not mislead to the application of linear regression models which would necessarily imply that the difference between “1” and “2” (e.g. “strong disagreement” and “weak disagreement”) is the same as the difference between “2” and “3” (e.g. “weak disagreement” and “weak agreement”) or “3” and “4” (e.g. “weak agreement” and “strong agreement”). In addition, linear regression models would imply that “2” is twice as high as the value “1” (e.g. persons with a “weak disagreement” would agree twice as strong to a political program as persons with a “strong disagreement”).

Continuous latent variable (which sometimes can be interpreted as varying utility) in ordered response models ($i = 1, \dots, n$):

$$y_i^* = \beta'x_i + \varepsilon_i$$

As in binary response models, $x_i = (x_{i1}, \dots, x_{ik})'$ is again a vector of k explanatory variables, $\beta = (\beta_1, \dots, \beta_k)'$ is the corresponding k -dimensional parameter vector, and ε_i is an error term. These unobservable latent variables can be related to the observed variables y_i or y_{ij} ($i = 1, \dots, n$; $j = 1, \dots, J$):

$$y_i = j \quad \text{if} \quad \kappa_{j-1} < y_i^* \leq \kappa_j$$

$$y_{ij} = \begin{cases} 1 & \text{if } y_i = j \\ 0 & \text{otherwise} \end{cases}$$

This threshold mechanism divides the latent variable y_i^* in J intervals by using $J + 1$ threshold parameters $\kappa_0, \kappa_1, \dots, \kappa_J$ with $\kappa_0 < \kappa_1 < \dots < \kappa_J$. According to this mechanism, higher values of the latent variable y_i^* lead to higher values of the ordered dependent variable y_i with the values or intervals $j = 1, \dots, J$. It follows:

$$y_i = 1 \quad \text{if} \quad \kappa_0 < y_i^* \leq \kappa_1 \quad \Leftrightarrow \quad \kappa_0 - \beta'x_i < \varepsilon_i \leq \kappa_1 - \beta'x_i$$

$$y_i = 2 \quad \text{if} \quad \kappa_1 < y_i^* \leq \kappa_2 \quad \Leftrightarrow \quad \kappa_1 - \beta'x_i < \varepsilon_i \leq \kappa_2 - \beta'x_i$$

$$y_i = 3 \quad \text{if} \quad \kappa_2 < y_i^* \leq \kappa_3 \quad \Leftrightarrow \quad \kappa_2 - \beta'x_i < \varepsilon_i \leq \kappa_3 - \beta'x_i$$

$$\vdots \qquad \qquad \qquad \vdots \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \vdots$$

$$y_i = J \quad \text{if} \quad \kappa_{J-1} < y_i^* \leq \kappa_J \quad \Leftrightarrow \quad \kappa_{J-1} - \beta'x_i < \varepsilon_i \leq \kappa_J - \beta'x_i$$

In order to comprise all real values of the latent variable y_i^* , it follows $\kappa_0 = -\infty$ and $\kappa_J = \infty$ so that the number of threshold parameters decreases to $J - 1$:

$$\begin{array}{lcl}
 y_i = 1 & \text{if } -\infty < y_i^* \leq \kappa_1 & \Leftrightarrow & -\infty < \varepsilon_i \leq \kappa_1 - \beta'x_i \\
 \vdots & \vdots & & \vdots \\
 y_i = J & \text{if } \kappa_{J-1} < y_i^* \leq \infty & \Leftrightarrow & \kappa_{J-1} - \beta'x_i < \varepsilon_i \leq \infty
 \end{array}$$

Remarks:

- The resulting $J - 1$ threshold parameters are required to divide the range of the unobservable latent variable y_i^* into J cells which correspond to the J categories of the observed ordered dependent variable y_i . Therefore, the categories of y_i represent a censored version of the true underlying values of the latent variable y_i^* (e.g. preferences)
- It follows that very low values of y_i^* are linked with $y_i = 1$ (e.g. “strong disagreement”), for $y_i^* > \kappa_1$ the value of y_i increases to 2 (e.g. “weak disagreement”), y_i increases to the value 3 for $y_i^* > \kappa_2$ (e.g. “weak agreement”) etc.
- In line with the discussion before, it can be seen that the difference between two levels of the ordered dependent variables y_i (e.g. “2” and “3” for “weak disagreement” and “weak agreement” or “3” and “4” for “weak agreement” and “strong agreement”) are not the same as on the scale of the latent variable y_i^* so that the threshold parameters capture a nonlinear transformation (and can be estimated within the ordered response models)

Different ordered response models result from different types of the density function of the latent variables y_i^* on the basis of a distribution assumption about ε_i with the distribution function $F_i(\varepsilon_i)$ as discussed later. If the $J - 1$ threshold parameters $\kappa_1, \kappa_2, \dots, \kappa_{J-1}$ and the parameters in β are summarized in the vector θ , it follows for the probability that y_i takes the value j ($j = 1, \dots, J$):

$$p_{ij}(x_i, \theta) = P(y_i = j | x_i, \theta) = P(y_{ij} = 1 | x_i, \theta) = F_i(\kappa_j - \beta'x_i) - F_i(\kappa_{j-1} - \beta'x_i)$$

On the basis of $\kappa_0 = -\infty$ and $\kappa_J = \infty$, it follows $F(-\infty) = 0$ and $F(\infty) = 1$. However, these probabilities comprise too many parameters so that not all threshold parameters are identified if a constant is included in the ordered response model. Therefore, one parameter has to be normalized. Common approaches are to set the first threshold parameter $\kappa_1 = 0$ or to drop the constant term from x_i . In the following, we consider the second approach.

Based on the aforementioned probabilities and the binary variables y_{ij} , the ML estimation of the $k + J - 1$ parameters β_1, \dots, β_k and $\kappa_1, \dots, \kappa_{J-1}$ in ordered response models (instead of $k \cdot (J-1)$ slope parameters and constants in pure multinomial logit and probit models) is identical to the ML estimation in multinomial discrete choice models. Therefore, it follows for the log-likelihood function:

$$\log L(\theta) = \sum_{i=1}^n \sum_{j=1}^J y_{ij} \log p_{ij}(x_i, \theta)$$

4.2 Ordered probit and logit models

Ordered probit models:

These ordered response models assume that the error terms ε_i are standard normally distributed (as in binary probit models which are special cases of ordered probit models with $J = 2$)

Choice probabilities in ordered probit models ($i = 1, \dots, n; j = 1, \dots, J$):

$$p_{ij}(x_i, \theta) = P(y_i = j | x_i, \theta) = \Phi_i(\kappa_j - \beta'x_i) - \Phi_i(\kappa_{j-1} - \beta'x_i)$$

For the specific probabilities this means:

$$P(y_i = 1 | x_i, \theta) = \Phi_i(\kappa_1 - \beta'x_i)$$

$$P(y_i = 2 | x_i, \theta) = \Phi_i(\kappa_2 - \beta'x_i) - \Phi_i(\kappa_1 - \beta'x_i)$$

$$P(y_i = 3 | x_i, \theta) = \Phi_i(\kappa_3 - \beta'x_i) - \Phi_i(\kappa_2 - \beta'x_i)$$

\vdots

$$P(y_i = J | x_i, \theta) = 1 - \Phi_i(\kappa_{J-1} - \beta'x_i)$$

→ As in binary probit models, the parameterization of the standard normal distribution of ε_i is not restrictive. In fact, the normalization of the normal distribution with an expected value of zero and variance one is necessary for the identification of the parameters in the choice probabilities

Ordered logit models:

These ordered response models are derived in the same way as ordered probit models and thus assume that the error terms ε_i are standard logistically distributed (as in binary logit models which are special cases of ordered logit models with $J = 2$)

Choice probabilities in ordered logit models ($i = 1, \dots, n; j = 1, \dots, J$):

$$p_{ij}(x_i, \theta) = P(y_i = j | x_i, \theta) = \Lambda_i(\kappa_j - \beta'x_i) - \Lambda_i(\kappa_{j-1} - \beta'x_i)$$

For the specific probabilities this means:

$$P(y_i = 1 | x_i, \theta) = \Lambda_i(\kappa_1 - \beta'x_i)$$

$$P(y_i = 2 | x_i, \theta) = \Lambda_i(\kappa_2 - \beta'x_i) - \Lambda_i(\kappa_1 - \beta'x_i)$$

$$P(y_i = 3 | x_i, \theta) = \Lambda_i(\kappa_3 - \beta'x_i) - \Lambda_i(\kappa_2 - \beta'x_i)$$

$$\vdots \quad \quad \quad \vdots$$

$$P(y_i = J | x_i, \theta) = 1 - \Lambda_i(\kappa_{J-1} - \beta'x_i)$$

→ In the same way as in the case of binary probit and logit models, the assumptions of standard normal or standard logistic distributions of ε_i in ordered probit or ordered logit models usually lead to very similar estimation results in practice across these two types of ordered response models (see later)

Similar to (pure) multinomial logit models, the ML estimators $\hat{\beta}_h$ can neither be interpreted as the estimators of the effect of the respective explanatory variable x_{ih} nor do they (generally, see later) indicate the direction of the estimator of marginal probability effects, i.e. a positive (negative) $\hat{\beta}_h$ does not necessarily lead to positive (negative) estimators of these effects. Instead, it follows for the estimator of the (partial) marginal probability effect of a (continuous) explanatory variable x_{ih} in general ordered response models as well as in ordered probit and logit models ($i = 1, \dots, n; j = 1, \dots, J$):

$$\frac{\partial \hat{p}_{ij}(x_i, \hat{\theta})}{\partial x_{ih}} = \left[f_i(\hat{\kappa}_{j-1} - \hat{\beta}'x_i) - f_i(\hat{\kappa}_j - \hat{\beta}'x_i) \right] \hat{\beta}_h$$

$$\frac{\partial \hat{p}_{ij}(x_i, \hat{\theta})}{\partial x_{ih}} = \left[\varphi_i(\hat{\kappa}_{j-1} - \hat{\beta}'x_i) - \varphi_i(\hat{\kappa}_j - \hat{\beta}'x_i) \right] \hat{\beta}_h$$

$$\frac{\partial \hat{p}_{ij}(x_i, \hat{\theta})}{\partial x_{ih}} = \left\{ \Lambda_i(\hat{\kappa}_{j-1} - \hat{\beta}'x_i) \left[1 - \Lambda_i(\hat{\kappa}_{j-1} - \hat{\beta}'x_i) \right] - \Lambda_i(\hat{\kappa}_j - \hat{\beta}'x_i) \left[1 - \Lambda_i(\hat{\kappa}_j - \hat{\beta}'x_i) \right] \right\} \hat{\beta}_h$$

In these equations $f_i(\cdot)$, $\varphi_i(\cdot)$, and $\Lambda_i(\cdot)[1-\Lambda_i(\cdot)]$ again symbolize general density functions of ε_i as well as the corresponding density functions of the standard normal and standard logistic distributions.

The estimators of a discrete change of $p_{ij}(x_i, \theta)$ due to a discrete change Δx_{ih} of a (particularly discrete) explanatory variable x_{ih} in general ordered response models as well as in ordered probit and logit models are ($i = 1, \dots, n; j = 1, \dots, J$):

$$\begin{aligned} \Delta \hat{p}_{ij}(x_i, \hat{\theta}) &= P(y_i = j | x_i + \Delta x_{ih}, \hat{\theta}) - P(y_i = j | x_i, \hat{\theta}) = \\ & \left[F_i(\kappa_j - \hat{\beta}'x_i - \hat{\beta}_h \Delta x_{ih}) - F_i(\kappa_{j-1} - \hat{\beta}'x_i - \hat{\beta}_h \Delta x_{ih}) \right] - \left[F_i(\kappa_j - \hat{\beta}'x_i) - F_i(\kappa_{j-1} - \hat{\beta}'x_i) \right] \\ \Delta \hat{p}_{ij}(x_i, \hat{\theta}) &= \left[\Phi_i(\kappa_j - \hat{\beta}'x_i - \hat{\beta}_h \Delta x_{ih}) - \Phi_i(\kappa_{j-1} - \hat{\beta}'x_i - \hat{\beta}_h \Delta x_{ih}) \right] - \\ & \left[\Phi_i(\kappa_j - \hat{\beta}'x_i) - \Phi_i(\kappa_{j-1} - \hat{\beta}'x_i) \right] \\ \Delta \hat{p}_{ij}(x_i, \hat{\theta}) &= \left[\Lambda_i(\kappa_j - \hat{\beta}'x_i - \hat{\beta}_h \Delta x_{ih}) - \Lambda_i(\kappa_{j-1} - \hat{\beta}'x_i - \hat{\beta}_h \Delta x_{ih}) \right] - \\ & \left[\Lambda_i(\kappa_j - \hat{\beta}'x_i) - \Lambda_i(\kappa_{j-1} - \hat{\beta}'x_i) \right] \end{aligned}$$

→ On the basis of the estimators of (partial) marginal probability effects and of discrete probability effects, it is again possible to estimate average marginal and discrete probability effects of an explanatory variable x_{ih} across all i as well as marginal and discrete probability effects of x_{ih} at the mean of the explanatory variables (the procedure for this estimation with STATA by considering differences of estimated probabilities is identical to the case of multinomial logit models)

Interpretation:

- The ML estimators $\hat{\beta}_h$ determine the estimated effect of the explanatory variable x_{ih} on the (indeed not interesting) latent variable y_i^* . Since the values of y_i^* are directly connected to the values of the ordered dependent variable y_i , the sign of $\hat{\beta}_h$ gives information about the direction of the estimated effect of x_{ih} on increasing values of y_i (but not on the single probabilities).
- Furthermore, the direction of the estimators of marginal and discrete probability effects for the categories $j = 1$ and $j = J$ is clear. For the estimators e.g. of marginal probability effects (with $\kappa_0 = -\infty$ and $\kappa_J = \infty$) it follows:

$$\frac{\partial \hat{p}_{i1}(x_i, \hat{\theta})}{\partial x_{ih}} = \left[f_i(-\infty) - f_i(\hat{\kappa}_1 - \hat{\beta}'x_i) \right] \hat{\beta}_h = -f_i(\hat{\kappa}_1 - \hat{\beta}'x_i) \hat{\beta}_h$$

$$\frac{\partial \hat{p}_{iJ}(x_i, \hat{\theta})}{\partial x_{ih}} = \left[f_i(\hat{\kappa}_{J-1} - \hat{\beta}'x_i) - f_i(\infty) \right] \hat{\beta}_h = f_i(\hat{\kappa}_{J-1} - \hat{\beta}'x_i) \hat{\beta}_h$$

Therefore, a positive (negative) $\hat{\beta}_h$ implies that an increasing x_{ih} leads to an increase (decrease) of $\hat{p}_{iJ}(x_i, \hat{\theta})$ and a decrease (increase) of $\hat{p}_{i1}(x_i, \hat{\theta})$.

- In contrast, the direction of the estimators of marginal and discrete probability effects for $j = 2, \dots, J - 1$ are ambiguous, even when positive (negative) $\hat{\beta}_h$ imply positive (negative) estimators of probability effects for high values of y_i and negative (positive) estimators for small values of y_i . The sign of the estimators of probability effects changes exactly once at different values of y_i ¹⁰ when moving from small to high values of y_i .

Example: Determinants of secondary school choice (I)

The determinants of the choice of 675 pupils in Germany between the three secondary school types Hauptschule, Realschule, and Gymnasium are again analyzed. In contrast to the previous application of a (pure) multinomial logit model, however, the natural rank order of the three categories is now used so that the coding of 1 for “Hauptschule”, the coding of 2 for “Realschule”, and the coding of 3 for “Gymnasium” of the dependent variable secondary school type (schooltype) indicates this order. The explanatory variables are the same as in the previous multinomial logit model analysis:

- Years of education of the mother (motheduc) as mainly interesting explanatory variable
- Dummy variable for labor force participation of the mother (mothinlf) that takes the value one if the mother is employed
- Logarithm of household income (loghhincome)
- Logarithm of household size (loghhsize)
- Rank by age among the siblings (birthorder)
- Year dummies for 1995-2002

The ML estimation of the ordered probit and logit models with STATA leads to the following results:

Example: Determinants of secondary school choice (II)

```
oprobit schooltype motheduc mothinlf loghhincome loghhsizе birthorder year1995 year1996
year1997 year1998 year1999 year2000 year2001 year2002
```

```
Ordered probit regression                               Number of obs   =           675
                                                         LR chi2(13)     =           203.31
                                                         Prob > chi2     =            0.0000
Log likelihood = -631.18653                             Pseudo R2       =            0.1387
```

schooltype	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	.2731466	.0279933	9.76	0.000	.2182808	.3280124
mothinlf	-.1649158	.0993593	-1.66	0.097	-.3596565	.0298249
loghhincome	.6492508	.1049847	6.18	0.000	.4434845	.8550172
loghhsizе	-.6145965	.2016538	-3.05	0.002	-1.009831	-.2193624
birthorder	-.1049289	.0576033	-1.82	0.069	-.2178293	.0079714
year1995	.0457511	.1877388	0.24	0.807	-.3222101	.4137124
year1996	.0898004	.1952066	0.46	0.645	-.2927974	.4723982
year1997	-.2803249	.1982527	-1.41	0.157	-.668893	.1082432
year1998	.0750879	.2120345	0.35	0.723	-.3404922	.4906679
year1999	-.1605177	.2046103	-0.78	0.433	-.5615465	.240511
year2000	.0009375	.2037127	0.00	0.996	-.3983321	.4002071
year2001	.0330622	.1972731	0.17	0.867	-.3535859	.4197104
year2002	-.105459	.1960686	-0.54	0.591	-.4897463	.2788283
/cut1	8.439737	1.061008			6.360201	10.51927
/cut2	9.369887	1.068849			7.274981	11.46479

Example: Determinants of secondary school choice (III)

```
ologit schooltype motheduc mothlnlf loghhincome loghhsizе birthorder year1995 year1996
year1997 year1998 year1999 year2000 year2001 year2002
```

```
Ordered logistic regression                               Number of obs   =           675
                                                         LR chi2(13)     =           205.38
                                                         Prob > chi2     =           0.0000
Log likelihood = -630.1549                               Pseudo R2      =           0.1401
```

schooltype	Coef.	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	.4678007	.0514731	9.09	0.000	.3669152	.5686862
mothlnlf	-.3025754	.1661858	-1.82	0.069	-.6282935	.0231427
loghhincome	1.149409	.1933411	5.94	0.000	.7704671	1.52835
loghhsizе	-1.063832	.3429954	-3.10	0.002	-1.736091	-.3915736
birthorder	-.1734154	.0975637	-1.78	0.075	-.3646368	.0178061
year1995	.0767835	.3175815	0.24	0.809	-.5456648	.6992318
year1996	.1352987	.3263589	0.41	0.678	-.504353	.7749504
year1997	-.4836479	.3314329	-1.46	0.144	-1.133245	.1659487
year1998	.1242217	.35564	0.35	0.727	-.5728199	.8212632
year1999	-.2312418	.3402744	-0.68	0.497	-.8981673	.4356837
year2000	-.0389321	.3385088	-0.12	0.908	-.7023971	.624533
year2001	.0343839	.3297169	0.10	0.917	-.6118494	.6806172
year2002	-.2241305	.3284821	-0.68	0.495	-.8679436	.4196826
/cut1	14.8607	1.967535			11.00441	18.717
/cut2	16.41517	1.987395			12.51994	20.31039

Example: Determinants of secondary school choice (IV)

Interpretation:

- The estimation results in the ordered probit and ordered logit models are qualitatively very similar
- The values of 203.31 and 205.39 of the likelihood ratio test statistic imply that the null hypothesis that all 13 parameters of the explanatory variables are zero (which would imply that the ordered response models only comprise the two thresholds) can be rejected at any common significance levels
- The parameter estimates for motheduc are positive and highly significantly different from zero due to the z statistics of 9.76 and 9.09 in the ordered probit and logit models, respectively
- These parameter estimates therefore imply that the years of education of the mother have a strong significantly positive effect on the choice of Gymnasium and a strong significantly negative effect on the choice of Hauptschule, whereas the effect on the choice of Realschule is ambiguous from these estimation results
- Similarly, loghincome has a strong significantly positive effect on the choice of Gymnasium, whereas loghsize has a strong significantly negative effect on the choice of Gymnasium

Example: Determinants of secondary school choice (V)

Wald and likelihood ratio tests:

As an example, the null hypothesis that neither `motheduc` nor `mothlnlf` has any effect on the secondary school choice, i.e. that the two corresponding parameters are zero, is tested after the ML estimation of the ordered probit model. The command for the Wald test in STATA is:

```
test motheduc mothlnlf

( 1)  [schooltype]motheduc = 0
( 2)  [schooltype]mothlnlf = 0

      chi2( 2) =    95.50
      Prob > chi2 =    0.0000
```

The corresponding commands for the likelihood ratio test in STATA are then:

```
estimates store unrestricted

oprobit schooltype loghhincome loghhsize birthorder year1995 year1996 year1997 year1998
year1999 year2000 year2001 year2002

estimates store restricted

lrtest unrestricted restricted

Likelihood-ratio test                    LR chi2(2) =    114.19
(Assumption: restricted nested in unrestricted)  Prob > chi2 =    0.0000
```

Example: Determinants of secondary school choice (VI)

The estimation of the average marginal probability effects of motheduc across all 675 pupils on the choice of Hauptschule, Realschule, and Gymnasium leads to the following (shortened) STATA results in the ordered probit model:

```
margins, dydx(motheduc) predict(outcome(1))
```

	Delta-method					
	dy/dx	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	-.0786704	.0075726	-10.39	0.000	-.0935124	-.0638283

```
margins, dydx(motheduc) predict(outcome(2))
```

	Delta-method					
	dy/dx	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	-.007464	.0025874	-2.88	0.004	-.0125352	-.0023928

```
margins, dydx(motheduc) predict(outcome(3))
```

	Delta-method					
	dy/dx	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	.0861344	.0073071	11.79	0.000	.0718128	.100456

Example: Determinants of secondary school choice (VII)

The estimation of corresponding marginal probability effects of motheduc at the means of the explanatory variables across all 675 pupils leads to the following (shortened) STATA results in the ordered probit model:

```
margins, dydx(motheduc) atmeans predict(outcome(1))
```

	Delta-method					
	dy/dx	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	-.0859161	.0086732	-9.91	0.000	-.1029152	-.0689171

```
margins, dydx(motheduc) atmeans predict(outcome(2))
```

	Delta-method					
	dy/dx	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	-.0199433	.0049997	-3.99	0.000	-.0297426	-.010144

```
margins, dydx(motheduc) atmeans predict(outcome(3))
```

	Delta-method					
	dy/dx	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	.1058594	.0110476	9.58	0.000	.0842065	.1275123

Example: Determinants of secondary school choice (VIII)

The estimation of the average marginal probability effects of motheduc across all 675 pupils on the choice of Hauptschule, Realschule, and Gymnasium leads to the following (shortened) STATA results in the ordered logit model:

```
margins, dydx(motheduc) predict(outcome(1))
```

	Delta-method					
	dy/dx	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	-.0795201	.0081198	-9.79	0.000	-.0954346	-.0636056

```
margins, dydx(motheduc) predict(outcome(2))
```

	Delta-method					
	dy/dx	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	-.0082689	.0030382	-2.72	0.006	-.0142235	-.0023142

```
margins, dydx(motheduc) predict(outcome(3))
```

	Delta-method					
	dy/dx	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	.087789	.0079214	11.08	0.000	.0722633	.1033146

Example: Determinants of secondary school choice (IX)

The estimation of corresponding marginal probability effects of motheduc at the means of the explanatory variables across all 675 pupils leads to the following (shortened) STATA results in the ordered logit model:

```
margins, dydx(motheduc) atmeans predict(outcome(1))
```

	Delta-method					
	dy/dx	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	-.0849093	.0089859	-9.45	0.000	-.1025214	-.0672972

```
margins, dydx(motheduc) atmeans predict(outcome(2))
```

	Delta-method					
	dy/dx	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	-.0276524	.007027	-3.94	0.000	-.0414252	-.0138797

```
margins, dydx(motheduc) atmeans predict(outcome(3))
```

	Delta-method					
	dy/dx	Std. Err.	z	P> z	[95% Conf. Interval]	
motheduc	.1125617	.0127688	8.82	0.000	.0875354	.1375881

Example: Determinants of secondary school choice (X)

Interpretation:

- The estimated average marginal probability effects or marginal probability effects at the means of the explanatory variables are very similar in the ordered probit and logit models, respectively, and strengthen the significantly positive effect of the years of education of the mother on the choice of Gymnasium and the significantly negative effect on the choice of Hauptschule
- The estimation results particularly imply that motheduc has a significantly negative effect on the choice of Realschule. This estimated negative effect is stronger on the basis of the marginal probability effects at the means of the explanatory variables than of the average marginal probability effects.
- The estimated average marginal probability effect of motheduc of -0.0075 in the ordered probit model implies that an increase of the years of education of the mother by one (unit) leads to an approximately estimated decrease of the choice probability for Realschule by 0.75 percentage points, whereas the estimated value of -0.0200 at the means of the explanatory variables implies an approximately estimated decrease by 2.00 percentage points
- While these estimated effects are very similar to those in the multinomial logit model, the estimated standard deviations are higher in the latter model which points to efficiency losses compared to ordered response models

Example: Determinants of secondary school choice (XI)

The estimation of the average probability of the choice of Hauptschule across all 675 pupils leads to the following STATA results in the ordered probit model:

```
margins, predict(outcome(1))
Predictive margins                    Number of obs   =           675
Model VCE      : OIM
Expression    : Pr(schooltype==1), predict(outcome(1))
```

	Margin	Delta-method Std. Err.	z	P> z	[95% Conf. Interval]	
---+-----						
_cons	.2931277	.015854	18.49	0.000	.2620545	.324201

The estimation of (average) discrete changes of probabilities due to a discrete change of an explanatory variable requires the estimation of (average) probabilities for specific values of the explanatory variable. For example, the average change of the probability of the choice of Hauptschule across all 675 pupils due to an increase of motheduc from the minimum value of seven years to the maximum value of 18 years of education can be estimated on the basis of the estimated average probabilities at these specific values. The corresponding STATA commands after the ML estimation of the ordered probit model are:

Example: Determinants of secondary school choice (XII)

```
margins, at(motheduc=7) predict(outcome(1))
```

```
Predictive margins          Number of obs   =          675
Model VCE      : OIM
Expression    : Pr(schooltype==1), predict(outcome(1))
at            : motheduc          =          7
```

	Margin	Delta-method Std. Err.	z	P> z	[95% Conf. Interval]	
---+---						
_cons	.6897138	.0423121	16.30	0.000	.6067836	.772644

```
margins, at(motheduc=18) predict(outcome(1))
```

```
Predictive margins          Number of obs   =          675
Model VCE      : OIM
Expression    : Pr(schooltype==1), predict(outcome(1))
at            : motheduc          =          18
```

	Margin	Delta-method Std. Err.	z	P> z	[95% Conf. Interval]	
---+---						
_cons	.0100681	.0052127	1.93	0.053	-.0001485	.0202848

The estimated average decrease of the probability of the choice of Hauptschule is therefore $0.6897 - 0.0101 = 0.6796$ or 67.96 percentage points for an increase from seven to 18 years

Example: Determinants of secondary school choice (XIII)

In contrast, the estimation of e.g. the probability of the choice of Hauptschule for the maximum value of motheduc = 18 years at the means of the other individual characteristics with STATA leads to the following results:

```
margins, at((means)_all motheduc=18) predict(outcome(1))
```

```
Adjusted predictions                               Number of obs   =           675
```

```
Model VCE      : OIM
```

```
Expression     : Pr(schooltype==1), predict(outcome(1))
```

```
at             : motheduc           =           18
                mothlnlf           =    .5525926 (mean)
                loghhincome        =   11.05839 (mean)
                loghhsize          =    1.412881 (mean)
                birthorder         =     1.76 (mean)
                year1995           =    .1377778 (mean)
                year1996           =     .12 (mean)
                year1997           =    .1111111 (mean)
                year1998           =    .0888889 (mean)
                year1999           =    .1007407 (mean)
                year2000           =    .1037037 (mean)
                year2001           =    .1185185 (mean)
                year2002           =    .117037 (mean)
```

		Delta-method				
	Margin	Std. Err.	z	P> z	[95% Conf. Interval]	
__cons	.0065595	.0037561	1.75	0.081	-.0008023	.0139213

4.3 Discussion of ordered probit and logit models

Underlying parallel regression assumption ($j = 1, \dots, J - 1$) with $F_i(\cdot) = \Phi_i(\cdot)$ in the ordered probit model and $F_i(\cdot) = \Lambda_i(\cdot)$ in the ordered logit model ($i = 1, \dots, n$):

$$P(y_i \leq j | x_i, \theta) = P(y_i^* \leq \kappa_j | x_i, \theta) = F_i(\kappa_j - \beta'x_i)$$

Consequences:

- These cumulative probabilities for several categories j only differ due to different threshold parameters, but not due to different parameter values
- For fixed cumulative probabilities the partial derivatives with respect to an arbitrary explanatory variable x_{ih} are identical for all j
- With this assumption, the following dummy variables can be specified:

$$d_{ij} = \begin{cases} 1 & \text{if } y_i \leq j \\ 0 & \text{if } y_i > j \end{cases}$$

As a consequence, the slope parameters, but not the threshold parameters, in ordered probit and logit models with two categories could be estimated by binary probit or logit models with these dummy variables as dependent variables.

- The underlying single index assumption is also the reason for the aforementioned property of ordered probit and logit models that the sign of the estimators of probability effects changes exactly once at different values of y_i when moving from small to high values of y_i 24

Latent variables in generalized ordered probit and logit models:

$$y_i^* = \beta_j' x_i + \varepsilon_i$$

Here it is thus allowed that the parameter vector β changes across j . It follows for the probabilities as discussed above with $F_i(\cdot) = \Phi_i(\cdot)$ in the generalized ordered probit model and $F_i(\cdot) = \Lambda_i(\cdot)$ in the generalized ordered logit model:

$$P(y_i \leq j | x_i, \theta) = F_i(\kappa_j - \beta_j' x_i)$$

These generalized ordered probit and logit models are clearly more flexible than conventional ordered probit and logit models:

- For example, the partial derivatives of fixed cumulative probabilities with respect to an arbitrary explanatory variable x_{ih} can vary across the categories j
- Furthermore, generalized ordered probit and logit models do not necessarily imply that the sign of the estimators of probability effects changes only once at different values of y_i when moving from small to high values of y_i

Statistical testing of conventional ordered probit and logit models:

- The null hypothesis of the simple ordered probit and logit models is that all parameter vectors β_j are identical across j (which implies the corresponding single index assumption)
- This hypothesis can e.g. be tested by using a likelihood ratio test when the generalized ordered probit or logit models are estimated (where the simple ordered probit or logit models are the restricted models)

Evaluation of the use of generalized ordered probit and logit models:

- These models do not ensure that the aforementioned probabilities are restricted to the interval between zero and one. Due to the possibly varying β_j , it is also possible that the estimated probabilities $P(y_i \leq j | x_i, \hat{\theta})$ decrease in j for specific values of the explanatory variables, which is, however, not logical.
- The ML estimators of the parameters (and thus e.g. estimators of marginal probability effects) in conventional ordered probit and logit models are inconsistent if the single index assumption is violated so that the ML estimation of generalized ordered probit or logit models is necessary in this case
- An alternative is the use of multinomial logit (or probit) models which also lead to consistent ML estimators of the parameters if the single index assumption is violated, although they are then inefficient
- To test the robustness of estimation results in empirical studies, it is certainly useful to compare the parameter estimates and the estimates of probability effects in different model approaches (e.g. binary probit and logit models, ordered probit and logit models, multinomial logit and probit models)

Interval data (e.g. income classes):

- For these data ordered probit and logit models can generally also be used
- The main difference to the previous analysis is that the thresholds values (i.e. the interval bounds such as income bounds) are known so that these threshold parameters need not to be estimated additionally